

MATH 271, HOMEWORK 9
DUE NOVEMBER 13TH

Problem 1. Compute the following:

(a)

$$[A] = \begin{pmatrix} 1 & 1 & 1 \end{pmatrix} \begin{pmatrix} 2 \\ 1 \\ 3 \end{pmatrix}.$$

(b)

$$[B] = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 5 & 6 & 7 & 8 \\ 9 & 10 & 11 & 12 \end{pmatrix} \begin{pmatrix} 3 & 2 \\ 2 & 3 \\ 3 & 2 \\ 2 & 3 \end{pmatrix}$$

(c) Take

$$[M] = \begin{pmatrix} 10 & 15 \\ 20 & 10 \end{pmatrix}$$

and

$$[N] = \begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix}.$$

Compute $[M][N]$ and $[N][M]$ to see that matrices do not commute in general.

Problem 2. A linear transformation $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ is given by the matrix

$$[T] = \begin{pmatrix} 1 & 2 & 0 \\ 2 & 1 & 2 \\ 0 & 2 & 1 \end{pmatrix}.$$

(a) Compute how T transforms the standard basis elements for \mathbb{R}^3 . That is, find

$$T(\hat{x}), \quad T(\hat{y}), \quad T(\hat{z})$$

and relate these values to the columns of $[T]$.

(b) Are the vectors $T(\hat{x})$, $T(\hat{y})$, and $T(\hat{z})$ linearly independent? Do these vectors form a basis for \mathbb{R}^3 ?

(c) If we apply this linear transformation to the unit cube (that is, all points who have (x, y, z) coordinates with $0 \leq x \leq 1$, $0 \leq y \leq 1$, and $0 \leq z \leq 1$), what will the volume of the transformed cube be? (*Hint: the determinant of this matrix $[T]$ provides us this information.*)

Problem 3. Consider some linear transformation $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$. Let $\vec{v}_1, \dots, \vec{v}_k$ be vectors in $\text{Null}(T)$.

- (a) Show that the span of these vectors is also in the nullspace of T .
- (b) How many linearly independent vectors can be in the nullspace?

Problem 4.

- (a) Show that for any 2×2 -matrix that the sign of the determinant changes if either a row or column is swapped. *Note: this is true for square matrices of any size.*
- (b) Show that for any 2×2 -matrix that multiplying a column by a constant scales the determinant by that constant as well. *Note: this is true for square matrices of any size.*
- (c) Show that for any 2×2 -matrix that adding a scalar multiple one column to the other will not change the determinant. *Note: this is true in broader generality. In fact, adding linear combinations of columns to another column will not change the determinant.*
- (d) Using these facts, argue why a square matrix with columns that are linearly dependent must have a determinant of zero.

Problem 5. Consider the equation

$$[A]\vec{v} = \vec{0},$$

where

$$[A] = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}.$$

- (a) Are the columns of $[A]$ linearly independent or dependent? Explain.
- (b) What vector(s) \vec{v} satisfy this equation? In other words, what is $\text{Null}([A])$?
- (c) Using what you found above, what must $\det([A])$ be equal to? *Hint: you do not need to compute the determinant!*

Problem 6. Compute the following determinants:

(a)

$$\det([A]) = \begin{vmatrix} -3 & 1 & 5 \\ -3 & 4 & 2 \\ -3 & 2 & 1 \end{vmatrix}$$

(b)

$$\det([B]) = \begin{vmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{vmatrix}$$

- (c) Compute $\det([A][B])$ using properties of the determinant. *Hint: this should be very quick to do. Do not compute the product of the matrices $[A]$ and $[B]$!*

Problem 7.

- (a) What does a zero determinant indicate about the solutions of a non-homogeneous system of linear equations? (Think geometrically!)
- (b) What does a zero determinant indicate about the solutions of a homogeneous system of linear equations? (Think geometrically!)

Problem 8. Given the matrices

$$[A] = \begin{pmatrix} 1 & 0 & 2 \\ 2 & 1 & 3 \\ -2 & -2 & 0 \end{pmatrix} \quad \text{and} \quad [B] = \begin{pmatrix} -3 & 1 & 1 \\ 2 & -2 & 4 \\ -1 & -1 & -1 \end{pmatrix}.$$

- (a) Compute $\text{tr}([A])$ and $\text{tr}([B])$.
- (b) Compute $\text{tr}([A][B])$ and compare it to $\text{tr}([B][A])$.