

MATH 272, HOMEWORK 10  
DUE APRIL 27<sup>TH</sup>

**Problem 1.** Consider the 1-dimensional wave equation given by

$$\left(-\frac{\partial^2}{\partial x^2} + \frac{1}{c^2} \frac{\partial^2}{\partial t^2}\right) u(x, t) = 0.$$

We'll consider two distinct scenarios. First, we'll take an infinitely long elastic rod and second we'll take a rod of finite length with Dirichlet boundary conditions.

(a) For a rod of infinite length, consider the initial conditions

$$u(x, 0) = \begin{cases} x + 1 & -1 \leq x \leq 0 \\ 1 - x & 0 \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases} \quad \text{and} \quad \frac{\partial}{\partial t} u(x, 0) = 0.$$

Find and plot the portion of the wave that moves to the right with  $c = 1$ .

(b) Let  $u_R(x, t)$  be your solution from (a), show that this satisfies the right-moving wave equation

$$\left(\frac{\partial}{\partial x} + \frac{1}{c} \frac{\partial}{\partial t}\right) u_R(x, t) = 0.$$

(c) Why is it that we can ignore the points where your function  $u_R(x, t)$  is not differentiable even though we are considering this as a solution to a PDE?

(d) For an elastic rod  $\Omega$  of finite length,  $\Omega = [0, 1]$ , assume that we take the Dirichlet conditions  $u(0, t) = 0 = u(1, t)$ . With the initial conditions

$$u(x, 0) = \sin(\pi x) \quad \text{and} \quad \frac{\partial}{\partial t} u(x, 0) = 0,$$

find the solution using d'Alembert's formula.

(e) Let  $w(x, t)$  be your solution for (d), show that it is indeed equal to

$$w(x, t) = \sin(\pi x) \cos(\pi ct).$$

(f) With your result from (e), explain how we can decompose a standing wave into a linear combination of two waves; one moving towards the left and one moving towards the right and both reflecting off the boundary.

**Problem 2.** Previously we studied the time-independent Schrödinger equation. Now, we can take a look at the time-dependent version given by

$$H\Psi(x, t) = i\hbar \frac{\partial}{\partial t} \Psi(x, t),$$

where  $H$  is the Hamiltonian operator. Consider the situation for the free particle in the 1-dimensional box of length  $L$  so that  $V(x) = 0$  and  $\Psi(0, t) = 0 = \Psi(L, t)$ .

- (a) Take a separation of variables ansatz and find a set of solutions (one for every positive integer  $n$ ) to the time-dependent equation.
- (b) Show that a super position of solutions is also a solution.
- (c) For a single state  $\psi_n(x, t)$ , show that

$$\int_0^L |\psi_n(x, t)|^2 dx,$$

is independent of  $t$ . This shows that the states  $\psi_n$  are *stationary* since their total probability does not depend on time.

**Problem 3.** Maxwell's equations are given as

$$\begin{aligned} \vec{\nabla} \cdot \vec{B} &= 0 & \vec{\nabla} \cdot \vec{E} &= \frac{\rho}{\epsilon_0} \\ \vec{\nabla} \times \vec{B} - \mu_0 \epsilon_0 \frac{\partial \vec{E}}{\partial t} &= \mu_0 \vec{J} & \vec{\nabla} \times \vec{E} + \frac{\partial \vec{B}}{\partial t} &= \vec{0} \end{aligned}$$

- (a) Look up each of the terms in the equations above and describe them.
- (b) Describe what each equation is saying and why these are PDEs.
- (c) In the absence of all charges we will have  $\vec{J} = \vec{0}$  and  $\rho = 0$ . Using that and the following two facts

$$\vec{\Delta} \vec{V} = \vec{\nabla}(\vec{\nabla} \cdot \vec{V}) - \vec{\nabla} \times (\vec{\nabla} \times \vec{V}) \quad \text{and} \quad \vec{\nabla} \times \frac{\partial \vec{V}}{\partial t} = \frac{\partial}{\partial t}(\vec{\nabla} \times \vec{V}),$$

derive the vector wave equations for light

$$\left( -\vec{\Delta} + \mu_0 \epsilon_0 \frac{\partial^2}{\partial t^2} \right) \vec{E} = \vec{0}$$

and

$$\left( -\vec{\Delta} + \mu_0 \epsilon_0 \frac{\partial^2}{\partial t^2} \right) \vec{B} = \vec{0}$$

- (d) From the equations you derived, determine the wave speed of light in the vacuum,  $c_0$ .